

Compact composition operators on H^2 and Hardy-Orlicz spaces

*Pascal Lefèvre, Daniel Li,
Hervé Queffélec, Luis Rodríguez-Piazza*

June 26, 2008

Abstract. *We compare the compactness of composition operators on H^2 and on Orlicz-Hardy spaces H^Ψ . We show in particular that there exists an Orlicz function Ψ such that $H^{3+\varepsilon} \subseteq H^\Psi \subseteq H^3$ for every $\varepsilon > 0$, and a composition operator C_ϕ which is compact on H^3 and on $H^{3+\varepsilon}$, but not compact on H^Ψ .*

Mathematics Subject Classification. Primary: 47B33; 47B10 – Secondary: 30C80

Key-words. Carleson function – Carleson measure – composition operator – Hardy spaces – Hardy-Orlicz spaces – Schatten classes

1 Introduction

Composition operators on the Hardy space H^2 have given rise to a lot of papers since the beginning of the seventies (see [11] and [2], and references therein, for an overview until the middle of the eighties). In particular, criteria of compactness ([12], [10]) or of membership in Schatten classes ([8] [9]) were found. But only few concrete examples of such operators were known (see [13], [1], [4], [14]). In [7], we constructed explicit examples of compact composition operators $C_\phi: H^2 \rightarrow H^2$ which are, or not, in some Schatten classes S_p . On the other hand, we study in [6] composition operators on Hardy-Orlicz spaces H^Ψ , and characterize their compactness. In this paper, we shall continue our investigation of composition operators on Hardy-Orlicz spaces, and study for which Orlicz functions Ψ the explicit examples of [7] are compact.

Let us describe more precisely the content of this paper.

In Section 3 and Section 4, we consider the Hardy-Orlicz space H^Ψ besides the Hardy space H^2 , and we prove that for every Orlicz function Ψ which does not satisfy the condition Δ_2 , there exists a symbol ϕ such that C_ϕ is compact on H^2 , but not compact on H^Ψ (Theorem 3.1). When the Orlicz function Ψ grows fast enough, we can ensure, beyond the non compactness of C_ϕ on H^Ψ , that, for every $p > 2$, $C_\phi: H^2 \rightarrow H^2$ is moreover in the Schatten class S_p (Proposition 3.4). But if Ψ does not grow too fast, this is not possible: if $C_\phi \in S_p$

for some $p > 0$, we must have the compactness of C_ϕ on H^Ψ (Proposition 3.3). When Ψ grows very fast, *i.e.* satisfies the condition Δ^2 , we show that C_ϕ may be in S_p for every $p > 0$, although C_ϕ is not compact on H^Ψ (Proposition 4.1); conversely, if C_ϕ is compact on H^Ψ , then C_ϕ must be in S_p for every $p > 0$ (Proposition 4.2).

In Section 5, we then characterize (Theorem 5.1) the Orlicz functions Ψ for which the composition operator C_{ϕ_θ} , which appeared in [7], Section 5, is compact on H^Ψ . When Ψ grows fast enough (in particular if Ψ satisfies the so-called condition Δ^1), then C_{ϕ_θ} is not compact on H^Ψ (Theorem 5.2); this gives, in particular, an improvement of Proposition 3.4. This characterization allows us to get the following striking result (Theorem 5.3): there exists an Orlicz function Ψ such that $H^{3+\varepsilon} \subseteq H^\Psi \subseteq H^3$ for every $\varepsilon > 0$, and a composition operator C_ϕ which is compact on H^3 and on $H^{3+\varepsilon}$, but not compact on H^Ψ .

2 Notation

We shall denote by $\mathbb{D}$ the open unit disk of the complex plane: $\mathbb{D} = \{z \in \mathbb{C}; |z| < 1\}$, and by $\mathbb{T} = \partial\mathbb{D}$ its boundary: $\mathbb{T} = \{z \in \mathbb{C}; |z| = 1\}$. We shall denote by m the normalized Lebesgue measure on $\mathbb{T}$.

For every analytic self-map $\phi: \mathbb{D} \rightarrow \mathbb{D}$, the composition operator C_ϕ is the map $f \mapsto f \circ \phi$. By Littlewood's subordination principle (see [3], Theorem 1.7), every composition operator maps continuously every Hardy space H^p ($p > 0$) into itself, as well as ([6], Proposition 3.12) the Hardy-Orlicz spaces H^Ψ (see below for their definition).

For every $\xi \in \mathbb{T}$ and $0 < h < 1$, the Carleson window $W(\xi, h)$ is the set

$$W(\xi, h) = \{z \in \mathbb{D}; |z| \geq 1 - h \text{ and } |\arg(z\bar{\xi})| \leq h\}.$$

For every finite positive measure μ on $\mathbb{D}$, one sets:

$$\rho_\mu(h) = \sup_{\xi \in \mathbb{T}} \mu[W(\xi, h)].$$

We shall call this function ρ_μ the *Carleson function* of μ .

When $\phi: \mathbb{D} \rightarrow \mathbb{D}$ is an analytic self-map of $\mathbb{D}$, and $\mu = m_\phi$ is the pullback measure defined on $\mathbb{D}$, for every Borel set $B \subseteq \mathbb{D}$, by:

$$m_\phi(B) = m(\{\xi \in \mathbb{T}; \phi^*(\xi) \in B\}),$$

where ϕ^* is the boundary values function of ϕ , we shall denote ρ_{m_ϕ} by ρ_ϕ . In this case, we shall say that ρ_ϕ is the Carleson function of ϕ .

For $\alpha \geq 1$, we shall say that μ is an α -Carleson measure if $\rho_\mu(h) \lesssim h^\alpha$. For $\alpha = 1$, μ is merely said to be a *Carleson measure*.

The Carleson Theorem (see [3], Theorem 9.3) asserts that, for $1 \leq p < \infty$ (actually, for $0 < p < \infty$), the canonical inclusion $j_\mu: H^p \rightarrow L^p(\mu)$ is bounded if and only if μ is a Carleson measure. Since every composition operator C_ϕ is

continuous on H^p , it defines a continuous map $j_\phi: H^p \rightarrow L^p(\mu_\phi)$; hence every pull-back measure μ_ϕ is a Carleson measure.

When $C_\phi: H^2 \rightarrow H^2$ is compact, one has, as it is easy to see:

$$(2.1) \quad |\phi^*| < 1 \quad a.e. \text{ on } \partial\mathbb{D}.$$

Hence, we shall only consider in this paper symbols ϕ for which (2.1) is satisfied (which is the case, as we said, when C_ϕ is compact on H^2). Recall ([6], Theorem 4.3) that, for every Orlicz function Ψ , as defined below, $C_\phi: H^2 \rightarrow H^2$ is compact whenever $C_\phi: H^\Psi \rightarrow H^\Psi$ is.

B. MacCluer ([10], Theorem 1.1) has shown, assuming condition (2.1), that C_ϕ is compact on H^p if and only if $\rho_\phi(h) = o(h)$, as h goes to 0.

Throughout this paper, the notation $f \approx g$ will mean that there are two constants $0 < c < C < +\infty$ such that $cf(t) \leq g(t) \leq Cf(t)$ (for t sufficiently near of a specified value), and the notation $f(t) \lesssim g(t)$, when t is in the neighbourhood of some value t_0 , will have the same meaning as $g = O(f)$.

Note that, in this paper, we shall not work, most often, with exact inequalities, but with inequalities up to constants. It follows that we shall not actually work with true Carleson windows $W(\xi, h)$, but with distorted Carleson windows:

$$\tilde{W}(\xi, h) = \{z \in \mathbb{D}; |z| \geq 1 - ah \text{ and } |\arg(z\bar{\xi})| \leq bh\},$$

where $a, b > 0$ are given constants. Since, for a given symbol ϕ , one has:

$$m_\phi(W(\xi, ch)) \leq m_\phi(\tilde{W}(\xi, h)) \leq m_\phi(W(\xi, Ch))$$

for some constants $c = c(a, b)$ and $C = C(a, b)$ which only depend on a and b , that will not matter for our purpose.

Let us now recall the definition of the Hardy-Orlicz spaces.

An Orlicz function is a non-decreasing convex function $\Psi: [0, \infty] \rightarrow [0, \infty]$ such that $\Psi(0) = 0$ and $\Psi(\infty) = \infty$. To avoid pathologies, we assume that the Orlicz function Ψ has the following additional properties: Ψ is continuous at 0, strictly convex (hence increasing), and such that

$$\frac{\Psi(x)}{x} \xrightarrow{x \rightarrow \infty} \infty.$$

This is essentially to exclude the case of $\Psi(x) = ax$.

The Orlicz space $L^\Psi(\mathbb{T})$ is the space of all (equivalence classes of) measurable functions $f: \mathbb{T} \rightarrow \mathbb{C}$ for which there is a constant $C > 0$ such that:

$$\int_{\mathbb{T}} \Psi\left(\frac{|f(t)|}{C}\right) dm(t) < +\infty$$

and then $\|f\|_\Psi$, the *Luxemburg norm*, is the infimum of all possible constants C such that this integral is ≤ 1 . The Hardy-Orlicz space H^Ψ is the space of all $f \in H^1$ such that the boundary values function f^* of f is in $L^\Psi(\mathbb{T})$. We refer to [6], Section 3 for more details.

3 Orlicz functions without condition Δ_2

In [6], Corollary 3.26, we showed that for every Orlicz function Ψ which satisfies the growth condition Δ^2 , there exists a symbol $\phi: \mathbb{D} \rightarrow \mathbb{D}$ which induces a compact composition on H^2 , but a non-compact composition operator on H^Ψ . We shall generalize this below. Recall that an Orlicz function Ψ satisfies the Δ_2 condition if there exists a constant $C > 0$ such that $\Psi(2x) \leq C\Psi(x)$ for x large enough.

Theorem 3.1 *For every Orlicz function Ψ which does not satisfy the condition Δ_2 , there exists an analytic self-map $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that C_ϕ is compact on H^2 but not compact on H^Ψ .*

Proof. $\Psi \notin \Delta_2$ means that $\limsup_{x \rightarrow \infty} \frac{\Psi(2x)}{\Psi(x)} = +\infty$. We can hence find a sequence (x_n) , which increases to $+\infty$, such that $\frac{\Psi(2x_n)}{\Psi(x_n)}$ increases to $+\infty$.

Set $h_n = \frac{1}{\Psi(x_n)}$ and $c_n = \frac{\Psi(x_n)}{\Psi(2x_n)}$. By construction, $(h_n)_n$ and $(c_n)_n$ decrease to 0. We may assume that $c_n < \pi$, $n \geq 1$.

We are going to construct an analytic function $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that:

$$(3.1) \quad \rho_\phi(h) = o(h),$$

but:

$$(3.2) \quad \rho_\phi(h_n) \geq c_n h_n, \quad \text{for all } n \geq 1.$$

Condition (3.1) will ensure that $C_\phi: H^2 \rightarrow H^2$ is compact, by MacCluer's theorem, and condition (3.2), which reads:

$$\rho_\phi(h_n) \geq \frac{1}{\Psi(2\Psi^{-1}(1/h_n))}, \quad \text{for all } n \geq 1,$$

will ensure (by [6], Theorem 4.11), that $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact.

For that purpose, we shall use the “general construction” made in [7], § 3.2. Let us recall this construction.

Let

$$(3.3) \quad f(t) = \sum_{k=0}^{\infty} a_k \cos(kt)$$

be an even, non-negative, 2π -periodic continuous function, vanishing at the origin: $f(0) = 0$, and such that:

$$(3.4) \quad f \text{ is strictly increasing on } [0, \pi].$$

The Hilbert transform (or conjugate function) $\mathcal{H}f$ of f is:

$$\mathcal{H}f(t) = \sum_{k=1}^{\infty} a_k \sin(kt).$$

We shall assume moreover that, as t tends to zero:

$$(3.5) \quad (\mathcal{H}f)'(t) = o(1/t^2).$$

Let now $F: \mathbb{D} \rightarrow \Pi^+ = \{\Re z > 0\}$ be the analytic function whose boundary values are:

$$(3.6) \quad F^*(e^{it}) = f(t) + i\mathcal{H}f(t).$$

One has:

$$F(z) = \sum_{k=0}^{\infty} a_k z^k, \quad |z| < 1,$$

and we define:

$$(3.7) \quad \Phi(z) = \exp(-F(z)), \quad z \in \mathbb{D}.$$

Since f is non-negative, one has:

$$\Re F(z) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) P_z(t) dt > 0,$$

P_z being the Poisson kernel at z , so that $|\Phi(z)| < 1$ for every $z \in \mathbb{D}$: Φ is an analytic self-map of $\mathbb{D}$, and $|\Phi^*| = \exp(-f) < 1$ a.e. . Note that the condition $f(0) = 0$ means that $\Phi^*(1) = 1$, so that $\|\Phi\|_{\infty} = 1$.

We then perturb Φ by considering:

$$(3.8) \quad M(z) = \exp\left(-\frac{1+z}{1-z}\right), \quad |z| < 1,$$

and

$$(3.9) \quad \phi(z) = M(z)\Phi(z), \quad |z| < 1.$$

We proved in [7], Lemma 3.6, that

$$(3.10) \quad hf^{-1}(h) \lesssim \rho_{\phi}(h) \lesssim hf^{-1}(2h)$$

when h tends to 0.

We shall now rely on the following simple lemma, whose proof is postponed.

Lemma 3.2 *Let (h_n) and (c_n) be two decreasing vanishing sequences of positive numbers, with $0 < h_n < \pi$, $0 < c_n < \pi$.*

There exists an even 2π -periodic $\mathcal{C}^2$ function $f: \mathbb{R} \rightarrow \mathbb{R}$ such that:

- 1) $f(0) = 0$ and f is strictly increasing on $[0, \pi]$;*
- 2) $f(c_n) \leq h_n$, for all $n \geq 1$.*

We are in the situation of the general construction, and by (3), we have:

$$\rho_\phi(h) = O(hf^{-1}(2h)) = o(h),$$

since $f^{-1}(2h) \rightarrow 0$ as $h \rightarrow 0$. MacCluer's criterion implies that $C_\phi: H^2 \rightarrow H^2$ is compact. On the other hand, (3) again implies that

$$\rho_\phi(h_n) \gtrsim h_n f^{-1}(h_n) \gtrsim c_n h_n,$$

by Lemma 3.2.

Therefore (3.2) holds, and this ends the proof of Theorem 3.1. $\square$

Remark. Actually, we do not use that f is $\mathcal{C}^2$, but only that it is continuous. However, if $f \in \mathcal{C}^2$, then $\widehat{f''} \in \ell_2(\mathbb{Z})$ and it follows, by the Cauchy-Schwarz inequality, since $\widehat{f''}(k) = (ik)^2 \widehat{f}(k)$, that $\sum_{k=0}^{\infty} k |\widehat{f}(k)| < +\infty$; hence both f and $\mathcal{H}f$ are $\mathcal{C}^1$ functions, and we proved in [7], Lemma 3.5, that then the Carleson function ρ_Φ of Φ is not $o(h)$ when h goes to 0, and so the composition operator $C_\Phi: H^2 \rightarrow H^2$ is not compact, although $|\Phi^*| = |\phi^*|$ on $\partial\mathbb{D}$ and $C_\phi: H^2 \rightarrow H^2$ is compact.

Proof of Lemma 3.2. Let $h_0 = c_0 = \pi$, and $\varphi: [0, \pi] \rightarrow \mathbb{R}_+$ be a step function such that:

$$(3.11) \quad \int_0^{c_n} \varphi(t) dt = h_n, \quad n = 1, 2, \dots$$

We can take $\varphi(t) = \frac{h_j - h_{j+1}}{c_j - c_{j+1}}$ for $t \in (c_{j+1}, c_j)$, $j = 0, 1, \dots$, since then:

$$\begin{aligned} \int_0^{c_n} \varphi(t) dt &= \sum_{j=n}^{\infty} \int_{c_{j+1}}^{c_j} \varphi(t) dt = \sum_{j=n}^{\infty} (c_j - c_{j+1}) \frac{h_j - h_{j+1}}{c_j - c_{j+1}} \\ &= \sum_{j=n}^{\infty} (h_j - h_{j+1}) = h_n. \end{aligned}$$

Let now $f: \mathbb{R} \rightarrow \mathbb{R}$ be the even 2π -periodic function whose values on $[0, \pi]$ are given by:

$$f(t) = \frac{1}{\pi^3} \int_0^t (t-u)^3 \varphi(u) du.$$

This function is clearly $\mathcal{C}^2$, with

$$f''(t) = \frac{6}{\pi^3} \int_0^t (t-u) \varphi(u) du,$$

and $f(0) = f'(0) = f''(0) = 0$; it is strictly increasing on $[0, \pi]$, and

$$f(c_n) = \frac{1}{\pi^3} \int_0^{c_n} (c_n - u)^3 \varphi(u) du \leq \frac{c_n^3}{\pi^3} \int_0^{c_n} \varphi(u) du \leq \int_0^{c_n} \varphi(u) du = h_n,$$

due to (3.11). Therefore f fulfills all the requirements of Lemma 3.2. $\square$

Remark. It is not clear whether one can ensure in Theorem 3.1 that the composition operator is moreover in some Schatten class S_p for $p < \infty$. It would be the case, by [7], Corollary 3.2, if one can construct a symbol ϕ such that its pull-back measure m_ϕ is α -Carleson for some $\alpha > 1$. This can be obtained if we can construct, in Lemma 3.2, our function f such that $f^{(N)}(0) \neq 0$ for some $N > 1$, since then $f(h) \approx h^N$ and $f^{-1}(h) \approx h^{1/N}$, and ϕ would be α -Carleson with $\alpha = 1 + \frac{1}{N}$, which would imply that $C_\phi \in S_p$ for $p > 2N$. However, in general, we cannot obtain that $f^{(N)}(0) \neq 0$ for some $N > 1$. Indeed, take $\Psi(x) = e^{(\log(x+1))^2} - 1$; then $h_n = \frac{1}{\Psi(x_n)} \approx \exp[-(\log x_n)^2]$, whereas, since

$$\Psi(2x) \approx \exp[(\log x)^2] \exp(2 \log x),$$

one has $c_n = \frac{\Psi(x_n)}{\Psi(2x_n)} \approx \exp(-2 \log x_n)$, so that $h_n = o(c_n^N)$ for every $N < \infty$. Since $f(c_n) \leq h_n$, all the possible derivatives of f at 0 must be zero.

Besides, when Ψ does not grow too fast, one has the following result.

Proposition 3.3 *Let Ψ be an Orlicz function such that, for every $A > 1$, one has, as x goes to infinity:*

$$(3.12) \quad \Psi(Ax) = O(\Psi(x)(\log \Psi(x))^\varepsilon), \quad \forall \varepsilon > 0.$$

Then, if $C_\phi \in S_p$ for some $p > 0$, the composition operator $C_\phi: H^\Psi \rightarrow H^\Psi$ is compact.

Remark. If the condition (3.12) is satisfied for some $A > 1$, it is actually satisfied for every $A > 1$. In fact, $\Psi(Ax) = O(\Psi(x)(\log \Psi(x))^\varepsilon)$ implies that $\Psi(A^2x) = O(\Psi(x)(\log \Psi(x))^{2\varepsilon})$, and we can replace A by $A^2, A^4, A^8, \dots$. Note also that we can replace the *big-oh* assumption by a *little-oh* one: if the proposition holds with a *little-oh* condition, it also holds with a *big-oh* one, because, for every $\varepsilon' > 0$, if one has (3.12), then, with $\varepsilon < \varepsilon'$:

$$\frac{\Psi(Ax)}{\Psi(x)(\log \Psi(x))^{\varepsilon'}} \leq K_\varepsilon (\log \Psi(x))^{\varepsilon - \varepsilon'} \xrightarrow{x \rightarrow \infty} 0.$$

Proof. If $C_\phi \in S_p$, then, by [7], Proposition 3.4, one has:

$$\rho_\phi(h) = o\left(\frac{h}{[\log(1/h)]^\delta}\right),$$

with $\delta = 2/p > 0$. Then, (3.12) with $x = \Psi^{-1}(1/h)$, gives:

$$\rho_\phi(h) = o\left(\frac{1}{\Psi(x)[\log \Psi(x)]^\delta}\right) = o\left(\frac{1}{\Psi(Ax)}\right) = o\left(\frac{1}{\Psi(A\Psi^{-1}(1/h))}\right),$$

which implies, by [6], Theorem 4.18, that C_ϕ is compact on H^Ψ . $\square$

For example, we have (3.12) if, for x large enough:

$$\Psi(x) = \exp(\log x \log \log \log x).$$

In fact, one has:

$$\frac{\Psi(Ax)}{\Psi(x)} = \exp \left[(\log x) \left(\log \frac{\log(\log x + \log A)}{\log \log x} \right) \right] \exp(\log A \log \log \log(Ax)),$$

and $\exp(\log A \log \log \log(Ax)) = (\log \log(Ax))^{\log A}$, whereas

$$\log(\log x + \log A) = \log \log x + \log \left(1 + \frac{\log A}{\log x} \right) \leq \log \log x + \frac{\log A}{\log x},$$

so:

$$\log \frac{\log(\log x + \log A)}{\log \log x} \leq \log \left(1 + \frac{\log A}{\log x \log \log x} \right) \leq \frac{\log A}{\log x \log \log x}.$$

Hence:

$$\frac{\Psi(Ax)}{\Psi(x)} \leq \exp \left(\frac{\log A}{\log \log x} \right) (\log \log(Ax))^{\log A} \lesssim (\log \log(Ax))^{\log A},$$

which gives (3.12) since, for every $\varepsilon > 0$:

$$(\log \log(Ax))^{\log A} = o((\log x)^\varepsilon) = o([\log \Psi(x)]^\varepsilon).$$

On the other hand, if we require that the Orlicz function Ψ grows sufficiently, we have:

Proposition 3.4 *Assume that the Orlicz function Ψ satisfies the condition:*

$$(3.13) \quad \limsup_{x \rightarrow \infty} \frac{\Psi(Ax)}{[\Psi(x)]^2} > 0,$$

for some $A > 1$.

Then, for every $p > 2$, there exists an analytic self-map $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that $C_\phi: H^2 \rightarrow H^2$ is in the Schatten class S_p , but $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact.

Note that condition (3.13) is not satisfied when $\Psi \in \Delta_2$, or for the Orlicz function $\Psi(x) = e^{(\log(x+1))^2} - 1$, i.e. $\Psi(x) \approx \exp((\log x)^2)$.

Proof. Condition (3.13) implies that there exist some $\delta > 0$ and a sequence of positive numbers x_n increasing to infinity such that:

$$\frac{\Psi(Ax_n)}{[\Psi(x_n)]^2} \geq \delta, \quad \text{for all } n \geq 1.$$

If one sets $h_n = \frac{1}{\Psi(x_n)}$, one has:

$$(3.14) \quad h_n^2 \geq \frac{\delta}{\Psi[A\Psi^{-1}(1/h_n)]}.$$

Now, take α such that $\frac{2}{p} + 1 < \alpha < 2$, and consider the symbol $\phi = \phi_2$ constructed in [7], Theorem 4.1. Its composition operator $C_\phi: H^2 \rightarrow H^2$ belongs to S_p ([7], Theorem 4.2). On the other hand, $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact, by [6], Theorem 4.18, since $\rho_\phi(h) \approx h^\alpha$ and hence $\rho_\phi(h_n)$ is not $o\left(\frac{1}{\Psi[A\Psi^{-1}(1/h_n)]}\right)$, by (3.14). $\square$

Condition (3.13) is a weaker growth condition than condition Δ^2 . Recall (see [6], § 2.1) that the Orlicz function Ψ satisfies the condition Δ^2 if, for some $A > 1$, one has $[\Psi(x)]^2 \leq \Psi(Ax)$ for x large enough. But condition Δ^2 can be re-stated as:

$$(3.15) \quad \liminf_{x \rightarrow \infty} \frac{\Psi(Ax)}{[\Psi(x)]^2} > 0.$$

Indeed, if one has $[\Psi(x)]^2 \leq \Psi(Ax)$ for x large enough, one obviously has $\liminf_{x \rightarrow \infty} \frac{\Psi(Ax)}{[\Psi(x)]^2} \geq 1$. Conversely, let $\delta = \liminf_{x \rightarrow \infty} \frac{\Psi(Ax)}{[\Psi(x)]^2}$. If $\delta > 1$, one clearly have $[\Psi(x)]^2 \leq \Psi(Ax)$ for x large enough. If $0 < \delta \leq 1$, then, for x large enough:

$$[\Psi(x)]^2 \leq \frac{2}{\delta} \Psi(Ax) \leq \Psi\left(\frac{2}{\delta} Ax\right),$$

by the convexity of Ψ , since $2/\delta \geq 1$. Hence $\Psi \in \Delta^2$ (with constant $2A/\delta$).

4 Orlicz functions with condition Δ^2

When $\Psi \in \Delta^2$, we have proved in [6], Corollary 3.26 a better result than the one in Proposition 3.4: *Whenever $\Psi \in \Delta^2$, there exists a symbol $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that the composition operator $C_\phi: H^2 \rightarrow H^2$ is Hilbert-Schmidt, whereas $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact.*

We have this improvement:

Proposition 4.1 *Let Ψ be an Orlicz function satisfying condition Δ^2 . Then, there exists a symbol ϕ such that $C_\phi: H^2 \rightarrow H^2$ belongs to S_p for every $p > 0$, but such that $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact.*

Proof. We shall use a *lens map*. One can find a definition of such maps in [11], page 27, but we shall use a slight variant of it: we shall use the same construction as in [7], Theorem 5.1 and Theorem 5.6, but with $f(z) = z^{1/2}$ instead of $f_\theta(z) = z(-\log z)^\theta$ or $f(z) = z \log(-\log z)$. We shall recall this construction at the beginning of the next section. Note that $\Re(re^{i\alpha})^{1/2} =$

$\sqrt{r} \cos(\alpha/2)$ is positive ($|\alpha| < \pi/2$), so the associate symbol ϕ maps $\mathbb{D}$ into itself.

As in the proof of [7], Theorem 5.1, we need only to consider $f(it)$, with $t > 0$. But $f(it) = \exp\left(\frac{1}{2} \log t + i\frac{\pi}{4}\right)$, so:

$$\Re f(it) = \sqrt{t} \cos(\pi/4) = \sqrt{t/2} \quad \text{and} \quad \Im f(it) = \sqrt{t} \sin(\pi/4) = \sqrt{t/2}.$$

Let us estimate $\rho_\phi(h)$. The condition $|\Re f(it)| \leq h$ implies $t \in [-2h^2, 2h^2]$, and, just using the modulus constraint, we get $\rho_\phi(h) \lesssim h^2$. For the converse, if $t \in [-2h^2, 2h^2]$, we have $|\Re f(it)| \leq h$ and $|\Im f(it)| \leq h$; then $\exp(f(it))$ belongs to a window centered at 1 and with size Ch . Hence $h^2 \lesssim \rho_\phi(h)$, and therefore $\rho_\phi(h) \approx h^2$.

It follows that $C_\phi: H^\Psi \rightarrow H^\Psi$ is not compact if $\Psi \in \Delta^2$. In fact, since $\Psi \in \Delta^2$, there is some $A_0 > 1$ such that $\Psi(A_0 x) \geq [\Psi(x)]^2$ for x large enough. We get, with $x = \Psi^{-1}(1/h)$:

$$\frac{1}{\Psi(A_0 \Psi^{-1}(1/h))} \leq h^2.$$

If C_ϕ were compact on H^Ψ , we should have ([6], Theorem 4.11):

$$\rho_\phi(h) = o\left(\frac{1}{\Psi(A_0 \Psi^{-1}(1/h))}\right),$$

which is not the case.

On the other hand, C_ϕ is in S_p for every $p > 0$. Indeed, $\phi(e^{it})$ is in the dyadic Carleson window:

$$W_{n,j} = \left\{ z \in \mathbb{D}; 1 - 2^{-n} \leq |z| < 1, \quad \frac{2j\pi}{2^n} \leq \arg(z) < \frac{2(j+1)\pi}{2^n} \right\},$$

($j = 0, 1, \dots, 2^n - 1$, $n = 1, 2, \dots$) if and only if, for $h_n = 2^{-n}$, one has, up to constants:

$$t \leq 2h_n^2 \quad \text{and} \quad j^2 h_n^2 \leq t \leq (j+1)^2 h_n^2.$$

This is possible for at most a fixed number, say N , of values of j ; hence:

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{j=0}^{2^n-1} 2^{np/2} m_\phi(W_{n,j})^{p/2} &\leq N \sum_{n=1}^{\infty} 2^{np/2} (\rho_\phi(2^{-n}))^{p/2} \\ &\leq N \sum_{n=1}^{\infty} 2^{np/2} 2^{-2np/2} = N \sum_{n=1}^{\infty} 2^{-np/2} < +\infty. \end{aligned}$$

Then Luecking's theorem ([8]) and [7], Proposition 3.3 imply that $C_\phi \in S_p$. $\square$

Now, in the opposite direction, one has proved in [6], Theorem 3.24, that if C_ϕ is compact on H^Ψ , with $\Psi \in \Delta^2$, then $C_\phi \in S_p(H^2)$ for every $p > 0$ (though we only stated that $C_\phi \in S_1$). We have the following improvement.

Proposition 4.2 *Let Ψ be an Orlicz function satisfying condition Δ^2 . Assume that C_ϕ is compact on H^Ψ . Then m_ϕ is an α -Carleson measure, for every $\alpha \geq 1$, and hence $C_\phi \in S_p$ for every $p > 0$.*

Proof. Since $\Psi \in \Delta^2$, there is some $A_0 > 1$ such that $\Psi(A_0 x) \geq [\Psi(x)]^2$, for x large enough. Hence, for every positive integer n , one has $\Psi(A_0^n x) \geq [\Psi(x)]^{2^n}$. Taking $x = \Psi^{-1}(1/h)$, one gets:

$$\Psi(A_0^n \Psi^{-1}(1/h)) \geq \frac{1}{h^{2^n}}.$$

Now, if C_ϕ is compact on H^Ψ , one has, by [6], Theorem 4.11, 1), with $A = A_0^n$:

$$\rho_\phi(h) \leq \frac{1}{\Psi(A_0^n \Psi^{-1}(1/h))};$$

therefore $\rho_\phi(h) \leq h^{2^n}$, and m_ϕ is a 2^n -Carleson measure.

The last assertion now follows from [7], Corollary 3.2. $\square$

5 Case of the symbols ϕ_θ

In [7], Section 5, we considered composition operators C_{ϕ_θ} which are, for every $\theta > 0$, compact on H^2 , and hence on H^p for every $p < \infty$. We are now going to examine when these composition operators are compact on H^Ψ .

Let us recall how the ϕ_θ 's are constructed.

For $\Re z > 0$, consider the principal determination of the logarithm $\log z$. For $\varepsilon > 0$ small enough, the function $f_\theta(z) = z(-\log z)^\theta$ has strictly positive real part on $V_\varepsilon = \{z \in \mathbb{C}; \Re z > 0 \text{ and } |z| < \varepsilon\}$ and $\Re f_\theta^*(z) > 0$ for all $z \in \partial V_\varepsilon \setminus \{0\}$. Now, if g_θ is the conformal mapping from $\mathbb{D}$ onto V_ε , which maps $\mathbb{T} = \partial \mathbb{D}$ onto ∂V_ε , and with $g_\theta(1) = 0$ and $g'_\theta(1) = -\varepsilon/4$, we set:

$$\phi_\theta = \exp(-f_\theta \circ g_\theta).$$

ϕ_θ maps $\mathbb{D}$ into itself and $|\phi_\theta^*| < 1$ on $\partial \mathbb{D} \setminus \{1\}$, and we proved ([7], Proposition 5.3) that:

$$(5.1) \quad \rho_{\phi_\theta}(h) \approx \frac{h}{(\log 1/h)^\theta}.$$

Now, recall ([6], Theorem 4.18) that C_{ϕ_θ} is compact on H^Ψ if and only if, for every $A > 0$:

$$\rho_{\phi_\theta}(h) = o\left(\frac{1}{\Psi[A\Psi^{-1}(1/h)]}\right), \quad h \rightarrow 0.$$

By (5.1), that gives, with $u = \Psi^{-1}(1/h)$:

Theorem 5.1 *The composition operator C_{ϕ_θ} is compact on H^Ψ if and only if for every $A > 1$, one has:*

$$(5.2) \quad \Psi(Au) = o\left(\Psi(u)[\log \Psi(u)]^\theta\right), \quad \text{as } u \rightarrow \infty.$$

For example, for every $\theta > 0$, C_{ϕ_θ} is compact on H^Ψ if $\Psi(x) \approx x^{\log \log \log x} = e^{(\log x)(\log \log \log x)}$, but is not compact on H^Ψ if $\Psi(x) \approx x^{\log \log x} = e^{(\log x)(\log \log x)}$.

Remark that this condition (5.2) is almost the same as condition (3.12): in (5.2) one is requiring the condition to be satisfied for some $\theta > 0$, whereas in (3.12) one is requiring it to be satisfied for every $\varepsilon > 0$ (see however the remark at the end of this Section). The *little-oh* assumption in (5.2) is actually not very important: suppose that, for every $A > 1$, one has $\Psi(Au) \leq C \Psi(u)[\log \Psi(u)]^\theta$ for $u \geq u_A$ (we need a uniform constant $C > 0$), then, for every $A > 1$, the convexity of Ψ gives, for every $\varepsilon > 0$, $\Psi(Au) \leq \Psi(A_\varepsilon u)$, with $A_\varepsilon = A/\varepsilon$; hence we get $\Psi(Au) \leq C \varepsilon \Psi(u)[\log \Psi(u)]^\theta$, and the *little-oh* property follows.

Note that, since C_{ϕ_θ} is in S_p for $p > 4/\theta$ ([7], Proposition 5.3), Theorem 5.1 is an improvement of Proposition 3.4, since (5.2) is not satisfied if and only if there exists some $A > 1$ such that $\limsup_{x \rightarrow \infty} \frac{\Psi(Ax)}{\Psi(x)[\log \Psi(x)]^\theta} > 0$, which is of course implied by condition (3.13):

Theorem 5.2 *Let $p > 0$ and $\theta > 4/p$. Assume that the Orlicz function Ψ satisfies, for some $A > 1$, the condition:*

$$(5.3) \quad \limsup_{x \rightarrow \infty} \frac{\Psi(Ax)}{\Psi(x)[\log \Psi(x)]^\theta} > 0.$$

Then the composition operator C_{ϕ_θ} , which is in $S_p(H^2)$, is not compact on H^Ψ .

If the Orlicz function Ψ satisfies condition Δ^1 , then condition (5.3) is satisfied.

Remark that, for $\theta > 2$, Theorem 5.2 improves Corollary 3.26 of [6]. Recall that the Orlicz function Ψ satisfies the condition Δ^1 if there is a constant $A > 1$ such that $x\Psi(x) \leq \Psi(Ax)$ for x large enough (see [6], § 2.1).

Proof of Theorem 5.2. The first part was explained before the statement. For the second part, assume that, for every $A > 1$, condition (5.2) is satisfied; in particular, we should have, for some $u_0 > 0$:

$$(5.4) \quad \Psi(Au) \leq \Psi(u)(\log \Psi(u))^\theta, \quad u \geq u_0.$$

Let

$$p_n = \Psi(A^n) \quad \text{and} \quad l_n = \log p_n.$$

One has then, by (5.4), for n large enough:

$$p_{n+1} \leq p_n(\log p_n)^\theta \quad \text{and} \quad l_{n+1} \leq l_n + \theta \log l_n.$$

Let

$$h(t) = \int_2^t \frac{dx}{\theta \log x}.$$

One has, since h' decreases:

$$0 < h(l_{n+1}) - h(l_n) \leq (l_{n+1} - l_n)h'(l_n) \leq (\theta \log l_n) \frac{1}{\theta \log l_n} \leq 1.$$

Hence $h(l_n) \lesssim n$, i.e.:

$$\int_2^{l_n} \frac{dx}{\theta \log x} \lesssim n.$$

Since this integral is $\gtrsim \frac{l_n}{\theta \log l_n}$, we get:

$$(5.5) \quad \frac{\log \Psi(A^n)}{\theta \log \log \Psi(A^n)} \lesssim n.$$

Hence:

$$\log \log \Psi(A^n) - \log \log \log \Psi(A^n) \lesssim \log n,$$

and so:

$$\limsup_{n \rightarrow \infty} \frac{\log \log \Psi(A^n)}{\log n} \leq 1.$$

On the other hand, $\Psi(A^n) \geq A^n$, i.e. $\log \Psi(A^n) \geq n \log A$.

Therefore $\log \log \Psi(A^n) \sim \log n$, and hence, using (5.5):

$$\log \Psi(A^n) \lesssim 2\theta n \log n.$$

But this prevents Ψ from satisfying the condition Δ^1 : if $\Psi \in \Delta^1$, there exists $A > 1$ such that $u\Psi(u) \leq \Psi(Au)$, for u large enough, and this implies, for n large enough, that $A^n \Psi(A^n) \leq \Psi(A^{n+1})$, and hence $A^{n(n+1)/2} \Psi(A) \lesssim \Psi(A^n)$, so that $\log \Psi(A^n) \gtrsim n^2$. $\square$

Remark. As a consequence of Theorem 5.1, one has the following striking result.

Corollary 5.3 *There exists an Orlicz function Ψ such that $H^{3+\varepsilon} \subseteq H^\Psi \subseteq H^3$ for every $\varepsilon > 0$, and a composition operator C_ϕ which is compact on H^3 and on $H^{3+\varepsilon}$, but not compact on H^Ψ .*

Remark. It is known ([13], Theorem 6.1), that the compactness of C_ϕ on H^{p_0} for some $p_0 < \infty$ implies its compactness on H^p for all $p < \infty$; this result follows from the Riesz factorization Theorem. It follows from Corollary 5.3 that there is no such factorization for Hardy-Orlicz spaces in general.

Proof. Consider the symbol $\phi = \phi_\theta$. The compactness of C_ϕ from H^2 into itself implies its compactness from H^p into itself for every $p \geq 1$; in particular its compactness on H^3 and on $H^{3+\varepsilon}$.

Let now Ψ the Orlicz function constructed in [5]. This function is such that $x^3/3 \leq \Psi(x)$ for all $x \geq 0$, but $\Psi(n!) \leq (n!)^3$ for all positive integer $n \geq 1$. Condition (5.2) is not satisfied when $\theta < 1$ because $\Psi(3.k!) \geq k.(k!)^3$ and

$$\Psi(k!)[\log \Psi(k!)]^\theta \leq (k!)^3 [3 \log(k!)]^\theta \lesssim (k!)^3 (k \log k)^\theta;$$

hence C_{ϕ_θ} is not compact on H^Ψ if $\theta < 1$. $\square$

Remark. Condition (5.2) depends on θ , but if Ψ is *regular*, the compactness of C_{ϕ_θ} on H^Ψ does not actually depend on θ , because, writing:

$$[\log \Psi(u)]^\theta \gtrsim \frac{\Psi(Au)}{\Psi(u)} = \frac{\Psi(Au)}{\Psi(\sqrt{Au})} \frac{\Psi(\sqrt{Au})}{\Psi(u)},$$

one gets:

$$\frac{\Psi(\sqrt{Au})}{\Psi(u)} \lesssim [\log \Psi(u)]^{\theta/2} \quad \text{or else} \quad \frac{\Psi(Au)}{\Psi(\sqrt{Au})} \lesssim [\log \Psi(u)]^{\theta/2}.$$

Hence, if Ψ satisfies the following condition of regularity:

$$(5.6) \quad \frac{\Psi(\sqrt{Au})}{\Psi(u)} \lesssim \frac{\Psi(Au)}{\Psi(\sqrt{Au})}, \quad \forall A > 1, \forall u > 0,$$

we get, for every $A > 1$ and every $u > 0$:

$$\frac{\Psi(\sqrt{Au})}{\Psi(u)} \lesssim [\log \Psi(u)]^{\theta/2},$$

and condition (5.2) remains true by replacing θ with $\theta/2$.

Note that condition (5.6) is satisfied whenever the Orlicz function Ψ satisfies the condition ∇_0 defined in [6], Definition 4.5; indeed, by Proposition 4.6 of [6], Ψ satisfies ∇_0 if and only if there exists some $u_0 > 0$ such that, for every $\beta > 1$, there exists $C_\beta \geq 1$ such that:

$$\frac{\Psi(\beta u)}{\Psi(u)} \leq \frac{\Psi(\beta C_\beta v)}{\Psi(v)}, \quad u_0 \leq u \leq v.$$

This condition implies (5.6), by taking $\beta = \sqrt{A}$ and $v = \sqrt{Au}$, since, by convexity, $\Psi(\beta C_\beta v) \leq C_\beta \Psi(\beta v)$.

When we impose in condition (5.6) an inequality with factor 1, for u large enough, one has:

$$[\Psi(\sqrt{Au})]^2 \leq \Psi(u) \Psi(Au), \quad \forall A > 1, \forall u \geq u_0,$$

and if one sets $u = e^x$ and $Au = e^y$, we get:

$$\log \Psi \left[\exp \left(\frac{x+y}{2} \right) \right] \leq \frac{1}{2} [\log \Psi(e^x) + \log \Psi(e^y)], \quad x_0 \leq x < y,$$

which means that the function $\kappa(x) = \log \Psi(e^x)$ is convex for x large enough. By [6], Proposition 4.7, that means that the Orlicz function Ψ satisfies the condition ∇_0 with constant 1. This regularity condition is satisfied, for example if $\Psi(x) = x^p$, $p \geq 1$, or $\Psi(x) = e^{(\log(x+1))^\alpha} - 1$, or else $\Psi(x) = e^{x^\alpha} - 1$, $\alpha \geq 1$.

6 Comparison with H^∞

It is well-known, and easy to see, that C_ϕ is compact on H^∞ if and only if $\|\phi\|_\infty < 1$. For composition operators, H^∞ is a very special case; indeed, one has:

Proposition 6.1

1) *Given any analytic self-map $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that $\|\phi\|_\infty = 1$, there exists an Orlicz function Ψ such that C_ϕ is not compact on H^Ψ .*

2) *Given any Orlicz function Ψ , there exists an analytic self-map $\phi: \mathbb{D} \rightarrow \mathbb{D}$ such that $\|\phi\|_\infty = 1$ and for which C_ϕ is compact on H^Ψ .*

Proof. 1) Since $\|\phi\|_\infty = 1$, one has $\rho_\phi(h) > 0$ for every $0 < h < 1$.

We construct piecewise a strictly convex function $\Psi: [0, +\infty[\rightarrow \mathbb{R}_+$ such that:

$$\Psi(2^{n+1}) \geq \frac{1}{\rho_\phi(1/\Psi(2^n))},$$

Then, if $h_n = 1/\Psi(2^n)$, one has $h_n \rightarrow 0$, and

$$\rho_\phi(h_n) \geq \frac{1}{\Psi(2\Psi^{-1}(1/h_n))}.$$

It follows from [6], Theorem 4.18, that C_ϕ is not compact on H^Ψ .

2) Let $a \in M^\Psi(\mathbb{T})$, the Morse-Transue space (see [6]) a positive function such that $a \notin L^\infty(\mathbb{T})$ and such that $\Psi \circ a \geq 2$, and let:

$$h = 1 - \frac{1}{\Psi \circ a}.$$

Since a is not essentially bounded, one has $\|h\|_\infty = 1$. Moreover, one has $h \geq 1/2$, so $\log h$ is integrable; we can hence define the outer function:

$$\phi(z) = \exp \left[\int_{\mathbb{T}} \frac{u+z}{u-z} \log h(u) dm(u) \right], \quad |z| < 1.$$

One has $|\phi^*(e^{i\theta})| = h(e^{i\theta})$, so ϕ is an analytic self-map from $\mathbb{D}$ into $\mathbb{D}$, and $\|\phi\|_\infty = 1$. Since, for every $A > 0$:

$$\Psi \left[A\Psi^{-1} \left(\frac{1}{1-|\phi^*|} \right) \right] = \Psi(Aa)$$

is integrable, because $a \in M^\Psi(\mathbb{T})$, the composition operator C_ϕ is M^Ψ -order bounded (see [6], Proposition 3.14), and hence is compact on H^Ψ . $\square$

References

- [1] T. Carroll and C. C. Cowen, Compact composition operators not in the Schatten classes, *J. Oper. Th.* 26 (1991), 109–120
- [2] C. C. Cowen and B. D. MacCluer, *Composition Operators on Spaces of Analytic Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL (1995).
- [3] P. Duren, *Theory of H^p Spaces*, Dover (2000)
- [4] M. Jones, Compact composition operators not in the Schatten classes, *Proc. Amer. Math. Soc.* 134 (2006), 1947–1953
- [5] P. Lefèvre, D. Li, H. Queffélec, and L. Rodríguez-Piazza, A criterion of weak compactness for operators on subspaces of Orlicz spaces, *to appear* in *Journal of Function Spaces and Applications*
- [6] P. Lefèvre, D. Li, H. Queffélec, and L. Rodríguez-Piazza, Composition operators on Hardy-Orlicz spaces, *preprint*, math.FA/0610905, *to appear* in *Memoirs Amer. Math. Soc.*
- [7] P. Lefèvre, D. Li, H. Queffélec, and L. Rodríguez-Piazza, Some examples of compact composition operators on H^2 , *to appear* in *Journal Funct. Anal.*
- [8] D. H. Luecking, Trace Ideal Criteria for Toeplitz Operators, *J. Funct. Anal.* 73 (1987), 345–368
- [9] D. H. Luecking and K. Zhu, Composition operators belonging to the Schatten ideals, *Amer. J. Math.* 114 (1992), 1127–1145
- [10] B. MacCluer, Compact composition operators on $H^p(B_N)$, *Michigan Math. J.* 32 (1985), 237–248
- [11] J. H. Shapiro, *Composition Operators and Classical Function Theory*, Universitext. Tracts in Mathematics, Springer-Verlag, New York (1993).
- [12] J. H. Shapiro, The essential norm of a composition operator, *Annals of Math.*, 125 (1987), 375–404
- [13] J. H. Shapiro and P. D. Taylor, Compact, Nuclear, and Hilbert-Schmidt Composition Operators On H^2 , *Indiana Univ. Math. J.* 23 (1973), 471–496
- [14] Y. Zhu, Geometric properties of composition operators belonging to Schatten classes, *Int. J. Math. Math. Sci.* 26 (2001), 239–248

Pascal Lefèvre, *Université d'Artois,*
Laboratoire de Mathématiques de Lens EA 2462,
Fédération CNRS Nord-Pas-de-Calais FR 2956,
Faculté des Sciences Jean Perrin,
Rue Jean Souvraz, S.P. 18,
62 307 LENS Cedex, FRANCE
pascal.lefevre@euler.univ-artois.fr

Daniel Li, *Université d'Artois,*
Laboratoire de Mathématiques de Lens EA 2462,
Fédération CNRS Nord-Pas-de-Calais FR 2956,
Faculté des Sciences Jean Perrin,
Rue Jean Souvraz, S.P. 18,
62 307 LENS Cedex, FRANCE
daniel.li@euler.univ-artois.fr

Hervé Queffélec, *Université des Sciences et Technologies de Lille,*
Laboratoire Paul Painlevé U.M.R. CNRS 8524,
U.F.R. de Mathématiques,
59 655 VILLENEUVE D'ASCQ Cedex, FRANCE
queff@math.univ-lille1.fr

Luis Rodríguez-Piazza, *Universidad de Sevilla,*
Facultad de Matemáticas, Departamento de Análisis Matemático,
Apartado de Correos 1160,
41 080 SEVILLA, SPAIN
piazza@us.es